

UNCLASSIFIED

MASTER COPY

FOR REPRODUCTION PURPOSES

2

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS DTIC FILE COPY	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S) ARO 21940.3-LS	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION U. S. Army Research Office	
6a. NAME OF PERFORMING ORGANIZATION University of California		6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211	
6c. ADDRESS (City, State, and ZIP Code) 405 Hilgard Avenue Los Angeles, CA 90024		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAA629-85-K-0113		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U. S. Army Research Office		8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
				WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Novel Approaches to the Study of Synaptic Function				
12. PERSONAL AUTHOR(S) Cameron B. Gundersen				
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 5/1985 TO 10/1988		14. DATE OF REPORT (Year, Month, Day) 16 January 1989
15. PAGE COUNT 3				
16. SUPPLEMENTARY NOTATION The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The final technical summary is presented on the accompanying pages:1&2.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL

DTIC  
ELECTE  
S FEB 17 1989 D  
D CB D

## Final Report

The original goal of the proposed experiments was to explore the possibility that mRNA-injected Xenopus oocytes could be used to reconstitute the transmitter release process. The hypothesis was that it might be possible to detect physiologically relevant acetylcholine release from Xenopus oocytes injected first with mRNA from cholinergic neurons and then with synaptic vesicles. For a number of technical reasons that have been discussed in prior reports, this approach has not been successful. In lieu of this effort, our primary focus has shifted toward attempts to clone the cDNA for two presynaptic proteins (the N-type calcium channel and the high affinity choline transporter) and one post-synaptic protein (a glutamate receptor). The status of these efforts is outlined below.

Because of this project's aim of reconstituting ACh release, we were using preparations of mRNA from the uniformly cholinergic neurons of Torpedo electric lobe. These mRNA preparations induce Xenopus oocytes to express functional choline acetyltransferase, an omega-conotoxin-sensitive calcium channel and a partially hemicholinium-3 inhibitable choline transporter. With the intention of trying to clone the cDNA for two of these proteins (the cDNA for choline acetyltransferase has been cloned), we fractionated mRNA on sucrose gradients and found that peaks of mRNA that induce choline uptake were around 2 Kb and that mRNA greater than 6 or 7 Kb induced expression of the calcium channel. In addition we observed that mRNA greater than 6 Kb in mass was predominantly responsible for the responses seen to kainic acid in oocytes injected with total electric lobe mRNA. Thus, in the course of screening size-selected cDNA libraries, we have tested for the presence of kainate receptors as well as calcium channels.

Our cloning strategy has been following several recent precedents in which mRNA is synthesized in vitro from DNA obtained from the cDNA library. This mRNA is injected into Xenopus oocytes which are then tested for the functional expression of the desired protein. The progress we have made for each of the three proteins we are trying to clone is summarized:

Choline transporter: A continuing problem for this project is that the increased uptake of choline in oocytes injected with Torpedo or rat brain mRNA has not been terribly sensitive to inhibition by hemicholinium-3, an agent that fully blocks high affinity choline transport in native tissue at about 1  $\mu$ M. Although we have performed some functional screening for this transporter using cDNA-library-derived mRNA, another approach potentially may be more useful. A Japanese group has cloned a choline transporter from yeast and we have found labelling of two bands of rat or Torpedo mRNA on Northern blots using the yeast cDNA as a probe. We are currently screening a rat cDNA library using the probe.

Calcium channel: To date we have not seen any evidence of calcium channel expression in oocytes injected with mRNA synthesized off of cDNA templates.

A-